

## A Study on the class of Semirings

M.Amala\*1, N.Sulochana2 and T.Vasanthi3

1,3Dept. of Applied Mathematics, Yogi Vemana University, Kadapa, Andhra Pradesh, India

2Asst. Prof., K.S.R.M College of Engineering, Kadapa, Andhra Pradesh, India

E-mail: amalamaduri@gmail.com, sulochananagam@gmail.com and vasanthitm@gmail.com

**Abstract :** In this paper, we study the class of Right regular and Multiplicatively subidempotent semirings. Especially we have focused on the additive identity 'e' which is also multiplicative identity in both semirings.

**Keywords:** Idempotent semiring, Multiplicatively subidempotent semiring, Periodic, positively totally ordered, Rectangular band.

### I. INTRODUCTION:

Various concepts of regularity on semigroups have been investigated by R.Croisot. His studies have been presented in the book of Clifford A.H. and G.B.Preston as R.Croisot theory one of the central places in the theory is held by the left regularity. K.S.S. Nambooripad studied on the structures of regular semigroups. we introduce the notion of Right regular semiring as a generalization of regular semiring. Sen, Ghosh & Mukhopadhyay studied the congruences on inverse semirings with the commutative additive reduct and Maity improved this to the regular semirings with the set of all additive idempotents a bi semilattice.

The study of regular semigroups has yielded many interesting results. These results have applications in other branches of algebra and analysis. Section one deals with introduction. Section two contains definitions. In third section we study on the class of right regular semiring. In last section we have given some results on ordered multiplicatively subidempotent semiring.

### II. PRELIMINARIES:

#### Definition 2.1:

A semiring S is a Right regular semiring, if S satisfies the identity  $a + xa + a = a$  for all a, x in S.

#### Definition 2.2:

A semigroup (S, +) is rectangular band if  $a = a + x + a$  for all a, x in S.

A semigroup (S, •) is rectangular band if  $a = axa$  for all a, x in S.

#### Definition 2.3:

An element a in a semigroup (S, +) is periodic if  $ma = na$  where m and n are positive integers. A semigroup (S, +) is periodic if every one of its elements is periodic.

#### Definition 2.4:

A semiring S is said to be idempotent if  $a + a = a$  and  $a^2 = a$  for all a in S.

#### Definition 2.5:

In a totally ordered semiring (S, +, •, ≤)

(i) (S, +, ≤) is positively totally ordered (p.t.o), if  $a + x ≥ a$ , x for all a, x in S

(ii) (S, •, ≤) is positively totally ordered (p.t.o), if  $ax ≥ a$ , x for all a, x in S.

### III. CLASSES OF RIGHT REGULAR SEMIRING:

In this section, the structures of Right regular semirings with different semiring properties are given. We have also framed examples in this chapter.

**Lemma 3.1:** Let S be a semiring which contains the additive and multiplicative identity 'e'. Then a ( $≠ e$ ) in S is a Right regular element if and only if  $xa = a$  for all x in S.

**Proof:** By hypothesis  $e$  is additive identity also multiplicative identity then  $a.e = e.a = a$  and  $e + a = a + e = a$  for all  $a, e$  in  $S$   
 Since  $a$  in  $S$  is a Right regular element then  $a + xa + a = a$  for all  $x$  in  $S$  which implies  $[e + x] a + a = a$   
 $\Rightarrow xa + a = a \Rightarrow xa = a$   
 Therefore  $xa = a$  for all  $x$  in  $S$   
 Now we have to prove that  $a \in S$  is a Right regular element  
 For this assume that  $xa = a$  for all  $x$  in  $S$   
 Let us consider  $a + xa + a$   
 The above equation can be written as  $[e + x] a + a = xa + a = xa = a$   
 Thus  $a + xa + a = a$  for all  $x$  in  $S$   
 Hence  $a$  in  $S$  is a Right regular element

**Theorem 3.2:** Let  $S$  be a semiring which contains an additive identity ' $e$ ' also multiplicative identity. If  $a$  ( $\neq e$ ) in  $S$  is a Right regular element, then  $a + x = a$  for all  $x$  in  $S$ .

**Proof:** Given that  $a$  in  $S$  is a Right regular element then  $a + xa + a = a$   
 By adding ' $x$ ' on both sides we get  $a + xa + a + x = a + x$   
 This implies  $a + xa + x = a + x \Rightarrow a + x(a + e) = a + x$   
 $\Rightarrow a + xa = a + x \Rightarrow xa = a + x$   
 Using above lemma  $xa = a$  the above equation reduces to the form  
 $a + x = a$  for all  $x$  in  $S$

**Proposition 3.3:** If  $S$  is a Right regular semiring with multiplicative identity ' $1$ ' and  $(S, \bullet)$  is a rectangular band, then  $(S, +)$  is periodic.

**Proof:** Since  $S$  is Right regular semiring then  $a + xa + a = a$  for all  $a, x$  in  $S$   
 This can also be written as  $a^2 + axa + a^2 = a^2$   
 Given that  $(S, \bullet)$  is rectangular band then  $axa = a$  for all  $a, x$  in  $S$   
 $\Rightarrow a.1.a + axa + a.1.a = a.1.a \Rightarrow a + a + a = a \Rightarrow 3a = a$   
 Thus  $(S, +)$  is periodic

#### IV. CLASSES OF MULTIPLICATIVELY SUBIDEMPOTENT SEMIRING:

In a semiring  $S$ , an element  $a$  is Multiplicatively Subidempotent if  $a + a^2 = a$ . A semiring  $S$  is Multiplicatively Subidempotent if and only if each of its elements is Multiplicatively Subidempotent. Multiplicatively Subidempotent semiring plays an important role in modal logic.

**Lemma 4.1:** Let  $S$  be a Multiplicatively Subidempotent semiring and ' $e$ ' be additive and multiplicative identity. Then  $S$  is an idempotent semiring.

**Proof:** By hypothesis  $e$  is an additive identity and also multiplicative identity then  $a.e = e.a = a$  and  $e + a = a + e = a$  for all  $a, e$  in  $S$   
 Since  $S$  is multiplicatively subidempotent  $a + a^2 = a$  for all  $a$  in  $S \rightarrow$  (1)  
 Equation (1) can be written as  $a(e + a) = a$  which implies  $a^2 = a \rightarrow$  (2)  
 Thus  $(S, \bullet)$  is a band  
 Adding  $a$  to both sides of equation (2) we obtain  $a + a^2 = a + a$   
 which implies  $a = a + a$  for all  $a$  in  $S \rightarrow$  (3)  
 From equations (2) and (3) we conclude that  $S$  is an idempotent semiring

**Example 4.2:** We have framed an example by considering the set  $S = \{a, x\}$  for above lemma which satisfies all the conditions of lemma.

+	a	x
a	a	a
x	a	x

.	a	x
a	a	a
x	a	x

**Theorem 4.3:** Let  $S$  be a totally ordered multiplicatively subidempotent semiring. If  $S$  contains multiplicative identity '1' in which additive identity 'e' is also multiplicative identity and  $(S, +)$  and  $(S, \cdot)$  are positively totally ordered then  $S$  is a mono semiring and the addition and multiplication are given by  $a + b = b + a = ab = ba = \max(a, b)$ .

**Proof:** By above lemma we have  $S$  is an idempotent semiring

Let  $a, b \in S$  and  $a < b$  implies  $a + a \leq a + b \leq b + b \Rightarrow a \leq a + b \leq b$

Since  $(S, +)$  is p.t.o this is possible only if  $a + b = b = \max(a, b)$

Also  $a < b$  implies  $a^2 \leq ab \leq b^2$  which implies  $a \leq ab \leq b$

Since  $(S, \cdot)$  is p.t.o this is possible only if  $ab = b = \max(a, b)$

**Example 4.4:** Here  $(S, +)$  and  $(S, \cdot)$  are p.t.o,  $y < x < a$  then  $a + b = b + a = ab = ba = \max(a, b)$ .

+	a	x	y
a	a	a	a
x	a	x	x
y	a	x	y

•	a	x	y
a	a	a	a
x	a	x	x
y	a	x	y

**Example 4.5:** Here  $(S, +)$  and  $(S, \cdot)$  are n.t.o,  $a < x < y$  then  $a + b = b + a = ab = ba = \min(a, b)$ .

+	a	x	y
a	a	x	y
x	x	x	y
y	y	y	y

•	a	x	y
a	a	x	y
x	x	x	y
y	y	y	y

**Note 4.6:**

(i) In the above theorem if  $(S, +)$  and  $(S, \cdot)$  are negatively totally ordered, then  $S$  is a mono semiring and the addition and multiplication are given by  $a + b = b + a = ab = ba = \min(a, b)$ .

(ii) In a t.o.s.r, if  $(S, +)$  is p.t.o and  $(S, \cdot)$  is n.t.o or vice-versa. We arrive contradiction to the hypothesis that additive identity 'e' is also multiplicative identity.

## REFERENCES

- [1.] Jonathan S.Golan, "The theory of Semirings with applications in Mathematics and Computer Science", Pitman Monographs and surveys in pure and applied mathematics, Vol – 54, Zongamann House, Burnt Mill, Harlow, Essex CM 20 JE. England: Longman Scientific and Technical, 1992.
- [2.] Jonathan S.Golan, "Semirings and their Applications", Kluwer Academic Publishers (1999).
- [3.] Jonathan S.Golan, "Semirings and Affine Equations over Them: Theory and Applications", Kluwer Academic Publishers (1999).
- [4.] Haixuan Yang and S.Ponizovskii, "Identities of Regular Semigroup Rings", Semigroup Forum Vol. 56 (1998) 293- 295 Springer-Verlag New York Inc.
- [5.] Kanchan Jana, "Quasi  $K$ -Ideals In  $K$ -Regular And Intra  $K$ -Regular Semirings", Pu. M.A.Vol. 22 (2011), No.1, Pp. 65-74.
- [6.] N.Kehayopulu, "On a characterization of regular duo le Semigroups", Maths. Balkonica 7 (1977), 181-186.
- [7.] K.S.S.Nambooripad, "Structure of regular semigroups I fundamental regular semigroups", Semigroup Forum, Vol.9 (1975), 354-363.
- [8.] M.Satyanarayana, "On the additive semigroup structure of semirings", Semigroup Forum, Vol.23 (1981), 7-14.
- [9.] M.Satyanarayana, "On the additive semigroup of ordered semirings". Semigroup Forum, 31 (1985), 193-199.
- [10.] M.K.Sen and S.K.Maity, "Semirings Embedded in a Completely Regular Semiring", Acta Univ. Palacki. Olomuc, Fac. rer. nat., Mathematica 43 (2004) 141–146.
- [11.] W.B.Vasanth Kandasamy, "Smarandache Semirings, Semifields and Semivector Spaces", Dept. of Mathematics, Indian Institute of Technology, Madras.